

Foot web pentosidine does not covary strongly with age in four species of wild seabirds

Angelika A. Aleksieva^a, Jason R. Treberg^{b,c}, Antony W. Diamond^d, Scott A. Hatch^e, Kyle H. Elliott^{a,*}

^a Department of Natural Resource Sciences, McGill University, Ste. Anne de Bellevue, Quebec H9X 3V9, Canada

^b Department of Biological Sciences, University of Manitoba, Winnipeg, Manitoba R3T 2N2, Canada

^c Centre on Aging, University of Manitoba, Winnipeg, Manitoba R3T 2N2, Canada

^d Atlantic Laboratory for Avian Research, University of New Brunswick, Fredericton, New Brunswick E3B 5A3, Canada

^e Institute for Seabird Research and Conservation, Anchorage, AK 99516, USA

ARTICLE INFO

Section Editor: Holly Brown-Borg

Keywords:

Advanced glycation endproducts

Age estimation

Biomarker

Senescence

ABSTRACT

Age is an important parameter for a variety of ecological applications, including population viability analyses, contaminants monitoring and targeting of individuals for conservation. While many organisms can be aged by annual rings, dentition and other techniques (i.e., fish otoliths, clam growth rings, mammal tooth wear), there are no minimally invasive biomarkers for accurately aging birds in the wild. For the past century, banding has been the only way to identify a bird of known age, which requires continuous effort on a large scale with possibly low return rates. Recent studies have identified pentosidine as a potential biomarker of chronological aging in several bird species. To test this idea in four species of long-lived seabirds, we collected skin biopsies from the foot webs of previously banded, known-age seabirds: black-legged kittiwakes (*Rissa tridactyla*; 0–19 y old), Atlantic puffins (*Fratercula arctica*; 5–26 y old), razorbills (*Alca torda*; 0–15 d old) and thick-billed murres (*Uria lomvia*; 0–35 y old). Foot web samples were specifically chosen because this was the least invasive site for substantial skin biopsy. Samples were analysed with high performance liquid chromatography to quantify pentosidine levels. Collagen levels were estimated through hydroxyproline assays to normalize pentosidine content across individuals. Kittiwakes displayed a weak correlation ($r^2 = 0.20$) between age and pentosidine/collagen. Puffins (adults only, $r^2 = 0.02$), razorbills (chicks only, $r^2 = 0.08$), and murres (adults, $r^2 = 0.04$) did not show any associations with age. We concluded that pentosidine content in the foot web does not appear to be a reliable method for aging seabirds in the wild. An absence of change in pentosidine in the foot web with age is further evidence that long-lived seabirds may maintain physiological performance into old age.

1. Introduction

Age is one of the most important parameters determining the health, reproductive potential and survival of organisms (Nussey et al., 2008; Ricklefs, 2008; Jones et al., 2014). Thus, knowledge of the age of an animal is critical for conservation applications, including population viability analyses, contaminants monitoring, and targeting of individuals for intervention (Fallon et al., 2006; Cooley, 2008; Holmes and Martin, 2009). While many organisms can be aged by annual rings, dentition and other techniques (i.e., fish otoliths, clam growth rings, mammal tooth wear), there are no biomarkers for accurately aging flying vertebrates in the wild beyond the first few years of life (Holmes and Martin, 2009; Herman et al., 2009; Munshi-South and Wilkinson,

2010; Jarman et al., 2015). Therefore marking a bird or bat during its hatch or birth year has been the only way to identify its exact age (Lincoln, 1921; Holmes and Martin, 2009; Munshi-South and Wilkinson, 2010). While osteologically-based age determination techniques are well-known for other vertebrates (Bagenal, 1974; Peabody, 1961; Reid, 1981), including many mammals (e.g., Morris, 1972; Grue and Jensen, 1979), none is well established for birds or bats (Broughton et al., 2002; Munshi-South and Wilkinson, 2010). Although the number of endosteal lamellae in transverse thin sections of the tibiotarsus showed some promise with small sample sizes in birds (Klomp and Furness, 1992), the technique was inaccurate with a larger number of individuals from one of the same species (Broughton et al., 2002). Moreover, such a procedure requires euthanizing the bird. It is thus

* Corresponding author.

E-mail address: kyle.elliott@mcgill.ca (K.H. Elliott).

<https://doi.org/10.1016/j.exger.2020.110833>

Received 15 August 2019; Received in revised form 2 January 2020; Accepted 6 January 2020

Available online 08 January 2020

0531-5565/ © 2020 Elsevier Inc. All rights reserved.

normally possible only to determine crudely whether birds are relatively young based on moult patterns, cranial pneumatization or morphometrics (e.g., McNeil and Burton, 1972; Sugimori et al., 1985; Siegel-Causey, 1989).

One reason why long-lived birds may be difficult to age in the wild is because they typically do not show the classic signs of physiological aging present in short-lived mammals. Actuarial senescence, the increasing rate of mortality with age, is more important in long-lived than short-lived wild animals as a higher proportion of mortality is attributable to senescence in the former relative to the latter (Ricklefs, 2008, 2010; Turbill and Ruf, 2010). Whereas relatively few individuals in short-lived species will live long enough to be senescent in the wild, a larger proportion of individuals in long-lived species will be senescent. Perhaps because of stronger selection for postponed physiological senescence in long-lived birds, studies of long-lived wild birds have detected few signs of declining immunity (Apanius and Nisbet, 2006; Lecomte et al., 2010), oxidative stress (Jimenez et al., 2019; Lecomte et al., 2010) or reproduction (Nisbet et al., 2002; Coulson and Fairweather, 2003; Lecomte et al., 2010) with age that routinely accompany physiological aging in mammals and short-lived birds (Cichon et al., 2003; Holmes and Ottinger, 2003; Saino et al., 2003; Moe et al., 2009; Palacios et al., 2007; these ideas are reviewed by Nisbet, 2001; Ricklefs, 2008, 2010; Holmes and Martin, 2009).

Aging is a complex phenomenon in which, at the cellular level, harmful changes can accumulate with time in each organism until its inevitable death. Biochemical changes at the level of cells and organelles ultimately result in organismal death (Harman, 1992, 2009; Martin et al., 1996). For example, the free radical theory of aging proposes that the generation of free radicals within the mitochondrial electron transport chain leads to progressive damage to organs, biological aging and, eventually, death (Harman, 1992; Wickens, 2001; Monaghan et al., 2009). Indeed, oxidative stress is speculated as a contributing factor in a number of aging-related diseases (Valko et al., 2007; Lushchak, 2014; Daffu et al., 2013). Another possible explanation of aging is the cross-linking theory of aging, where sugars modify body proteins through the Maillard reaction (Monnier and Cerami, 1981; Grandhee and Monnier, 1991; Iqbal et al., 1999a,b). The results of cross-linking are advanced glycation end products (AGEs) whose formation is accelerated by oxidative stress (Wickens, 2001). Advanced glycation endproducts can damage the organism directly, by modifying intracellular or extracellular proteins, and, indirectly, by binding to the receptor of advanced glycation end products (RAGE). Ligand binding to RAGE increases intracellular reactive oxygen species formation, and therefore leads to oxidative stress and could contribute to aging as described by the free radical theory (Daffu et al., 2013; Nowotny et al., 2015). One such AGE is pentosidine (Sell and Monier, 1989) and named after originally being synthesized from pentose—although pentosidine can be derived from other sugars. At the end of the Maillard reaction, skin pentosidine cross-links the arginine and lysine residues within collagen (Sell and Monnier, 1989; Sell et al., 1998; Dyer et al., 1991). Cross-linking alters collagen's structure and mechanical properties, which makes the protein less degradable by proteases (Nowotny et al., 2015). Furthermore, human neurons exposed to pentosidine increased their reactive oxygen species generation and apoptotic cell death (Nitti et al., 2005). Thus, AGEs could link biological age (physiological state) to chronological age (time since birth) (Sell et al., 2003; Chaney et al., 2003; Fallon et al., 2006). Indeed, due to its stable, fluorescent, and accumulating properties, previous studies confirmed pentosidine as potential biomarker of age in mammals, birds, and, most recently, in turtles (Sell et al., 1998; Cooney et al., 2010; Dammann et al., 2011; Iverson et al., 2017).

Pentosidine increases significantly with age in chickens (Iqbal et al. 1997, 1999a), California gulls (*Larus californicus*; Chaney et al., 2003), ruffed grouse (*Bonasa umbellus*; Fallon et al., 2006), monk parakeets (*Myiopsitta monachus*; Cooney et al., 2010) and double-crested cormorants (*Phalacrocorax auritus*; Fallon et al., 2006; Cooney, 2008; Dorr et al., 2017), but not common gulls (*Larus canus*; Rattiste et al., 2015), kakapos (*Strigops*

habroptilus; Campion, 2015) or bridled terns (*Onychoprion anaethetus*; Labbé et al., 2019). Thus, results so far are consistent with the idea that physiological senescence (as shown by pentosidine content increasing with age) is evident in short-lived but not long-lived birds, but few long-lived birds have been tested. In cormorants, pentosidine levels in the patagium were lower than breast skin, but both tissues showed a similar increase with age (Cooney et al., 2010). Levels in foot webs were also lower than in breast skin (Chaney et al., 2003). We examined levels in the foot web, to determine if biopsies from the foot web could be a non-lethal and minimally invasive technique for measuring age in waterbirds, as biopsies from the foot web may cause less damage and risk of infection than biopsies from the breast or patagium. We examined whether pentosidine can be used as a biomarker of chronological age in four species of long lived seabirds: black-legged kittiwakes (*Rissa tridactyla*), Atlantic puffins (*Fratercula arctica*), razorbills (*Alca torda*), and thick-billed murres (*Uria lomvia*). We expected pentosidine concentrations to increase with age among all individuals, with species-specific rates varying with maximum lifespan—as found in other taxa (Fallon et al., 2006).

2. Materials and methods

2.1. Tissue sampling

Four species of seabirds were sampled during their breeding season in 2015 and 2016 under McGill animal use protocol 2015-7599. Only birds banded as chicks, and thus of known ages, were targeted. Adults were trapped by various methods and put in cloth bags to minimize stress. Middleton Island, Alaska (59° 26' N, 146° 20' W), in the Gulf of Alaska (Gill and Hatch, 2002), is the site of an abandoned U.S. Air force defense radar tower that black-legged kittiwakes use for nesting. The buildings' sides were upgraded with wooden panels, nest ledges, and sliding panes of one-way mirror glass, thus creating a research habitat consistently of hundreds of nest sites easy to access and monitor. Since 1984, kittiwake adults and chicks have been banded on Middleton Island during every breeding season (Gill and Hatch, 2002). At Middleton, kittiwake reproductive success increases until about age 12, and then decreases rapidly (Elliott et al., 2014, with little change in muscle physiology Brown et al., 2019). In 2015, we collected samples from adults either nesting on or prospecting for sites on the tower, as well as from chicks hatched on it. Adults were captured opportunistically with wire hooks passed through slots in the wall at or below the level of their nests (Chivers et al., 2015). Chicks were monitored daily until 40 days since hatching, and were taken from their nest on the day of sampling.

Machias Seal Island, New Brunswick (44° 30' N, 67° 06' W) lies 18 km off the coast of Maine (Diamond and Devlin, 2003; Kelly et al., 2015). Since 1980 and 1995 respectively, Atlantic puffins and razorbills have been banded on Machias Seal Island (Kelly et al., 2015). In 2016, we collected samples from Atlantic puffins (adults and chicks) and razorbill chicks spread across the island. Puffins nest in burrows, and most adults were captured by extracting them directly from their nest sites. Occasional individuals were caught with drop-box traps or mist nets. While still in the egg stage, most study burrows were flagged and regularly checked until their hatch date. Each had an assigned sampling day and was not inspected until a few days before that date to minimize stress and abandonment by the parents (Rodway et al., 1996). Chicks younger than 36 days were caught in the burrow. Puffin chicks, older than 36 days, were caught at night during their first trip to the sea. Razorbills nest among boulders and adults are hard to catch, so we did not capture any adults. Razorbill chicks were monitored and sampled similarly to the puffin chicks at about age 10 d.

Coats Island, Nunavut (62° 57' N, 82° 45' W) is an island in northern Hudson Bay (Gaston et al., 1994). Thick-billed murres on Coats Island are located in two sub-colonies 1.5 km apart and have been banded since 1981 (Gaston et al., 1994; Elliott et al., 2015). At Coats, reproductive success increases until age 25, and then decreases slightly (Elliott et al., 2014). The oldest record for a thick-billed murre is 37

years (Jimenez et al., 2019), and our age range of 0–35 y is therefore representative of the age range present in the wild population. In 2016, adult murre were captured using a noose pole during the incubation period. Murre chicks were also caught by noosing near the top of the cliffs within 5 days of their hatch.

For all species, feet were cleaned with isopropanol, followed by puncturing an interdigital foot web with a 6 mm biopsy punch. The collected skin sample was stored in a labeled centrifuge tube. We collected one sample per bird, punching through the entire web. The foot hole was treated with antibiotic ointment to prevent infection and pressed with styptic powder in between two cotton balls until hemostasis. The full procedure lasted for a maximum of 10 min before releasing the bird. Samples were transported, surrounded by icepacks, from the field to a -20°C freezer within 5 to 60 min after being collected. From 3 weeks to 3 months later, they were transported with dry ice to our laboratory facilities. Sex was already known for black-legged kittiwakes based on copulation position and for thick-billed murre based on genetics.

2.2. Sample processing

Regardless of species, all samples were processed identically in a method modified from that described by Iqbal et al. (1997), Chaney et al. (2003), and Fallon et al. (2006). Biopsies were scraped clean, minced with razor blades and transferred to glass tubes. To remove lipids from skin biopsies, each tube was filled with 2 mL of a 2:1 chloroform/methanol solution, placed in a shaker, and left mixing at $4-6^{\circ}\text{C}$ for 18 h. We discarded the chloroform/methanol waste by pipetting it out of the tubes and leaving the rest to air-dry for 1 h. Once completely dry, the residue was rehydrated, hydrolyzed in 1 mL of 6 M HCl and heated to 110°C for 18 h. After cooling down, the samples were centrifuged to settle the suspended remains and retain only the liquid portion. The skin extracts were then neutralized by titration with NaOH. The resulting saline solutions were frozen at -80°C and transferred to a freeze dryer (Labconco 75040) to undergo sublimation. Finally the samples were reconstituted in 1 mL of Milli-Q water and filtrated with $0.22\ \mu\text{m}$ syringe filters. We examined the effect of reconstituting the sample in 0.25–1.5 mL of Milli-Q water, and there was no difference in the recovered pentosidine or collagen values among the different dilutions. Some spiked samples with pentosidine (from individuals not otherwise used in this study) were used to help identify peaks on HPLC chromatographs. The percent recovery of $0.080\ \text{pmol}/\mu\text{L}$ spiked pentosidine in five murre foot web samples was 99.7 ± 4.0 (SD) %.

2.3. Pentosidine measurement

Pentosidine (Cayman Chemicals, Ann Arbor, MI), in a crystalline solid state is stable for up to 2 years when kept at -20°C (we obtained standard curves varying by $< 5\%$ when comparing pentosidine kept frozen over that time period). Once diluted in water, the aqueous solution should not be stored for more than a day (standard curves of pentosidine decreased by about 0.8% per hour such that apparent levels were 20% of actual levels after 100 h). Therefore, the stock was divided in multiple aliquots and frozen until needed during the experiment. Before each experiment, one aliquot was thawed and further diluted into 6 standard concentrations of $0.5\ \text{nmol}/\text{L}$, $1.0\ \text{nmol}/\text{L}$, $2.0\ \text{nmol}/\text{L}$, $4.0\ \text{nmol}/\text{L}$, $8.0\ \text{nmol}/\text{L}$, and $16\ \text{nmol}/\text{L}$. Similarly to the sample extracts, each standard was filtered with $0.22\ \mu\text{m}$ syringe filters before its transfer to a glass vial.

Pentosidine was measured through modifications of the reverse phase high-performance liquid chromatography (HPLC) techniques previously described by Iqbal et al. (1997), Chaney et al. (2003), and Fallon et al. (2006). Pentosidine concentrations were estimated in duplicate using a Dionex UltiMate 3000 HPLC work station with an in-line fluorescence detector (UltiMate 3000, excitation 325 nm, emission 370 nm). The mobile phase consisted of acetonitrile, Milli-Q water and $0.01\ \text{M}$ heptafluorobutyric acid (HFBA). Injections ($100\ \mu\text{L}$) were

separated using a linear gradient of 5–30% acetonitrile, in water with HFBA from 0 to 35 min using a C8 column (Acclaim $4.6 \times 10\ \text{mm}$) and flowrate of $1\ \text{mL}/\text{min}$. The resulting elutions were visualized as chromatograms with peaks of different magnitudes at specific timespans. Integration of those peaks was performed in Chromeleon 7 version 7.1.2.1478 software package. A pentosidine standard was processed after every 4 to 5 skin extracts to confirm pentosidine's exact elution timing. The column was cleaned and equilibrated for a minimum of 20 and 15 min respectively between injections.

2.4. Collagen measurement

Assuming that 14% of collagen by weight is hydroxyproline, we used a spectrophotometric assay to measure each sample's hydroxyproline content using a commercial kit (MAK008, Sigma-Aldrich, Missouri, St Louis; Maekawa et al., 1970). Hydroxyproline was measured for all hydrolyzed skin extracts in duplicate (coefficient of variation averaged 8.9%) with a Fluostar omega microplate reader (emission wavelength of 560 nm). The results were examined with MARS Data Analysis version 2.41 software. The assay standards were diluted in water with 6 M NaCl to mimic the NaCl formed during sublimation of neutralized samples of hydrolyzed skin tissue; an initial attempt using KCl led to inaccurate data compared with water.

2.5. Statistical analysis

Statistical regression analyses using general linear models (LM) were carried out with R to determine the relationship between pentosidine or hydroxyproline concentrations and age + age² + sex. We considered an age² term to account for non-linear relationships. We included sex as a covariate only for murre and kittiwakes as they were the only two species where we had sex information. We used a backwards selection procedure where we included age, age² and sex as covariates and then dropped non-significant terms systematically based on their non-significance. We used a significance level of $P < 0.05$, and considered significant relationships with an $R^2 > 0.5$ as useful for age determination.

3. Results

3.1. Kittiwakes

We sampled 30 black-legged kittiwake adults aged 3–19 years, with most single age groups having two individuals per age. After normalizing across individuals using the hydroxyproline assay results, pentosidine concentrations in collagen ranged from $1.31\ \text{pmol}$ pentosidine/mg collagen to $8.16\ \text{pmol}/\text{mg}$ (Fig. 1). These concentrations increased linearly with age: pentosidine = $0.1689 * \text{age} + 2.5015$, $R^2 = 0.20$ ($t_{29} = 2.64$, $P = 0.01$; omnibus GLM age²: $t_{27} = 1.49$, $P = 0.15$; sex: $t_{27} = 0.42$, $P = 0.69$). Average pentosidine concentrations in each group increased linearly with age: pentosidine = $0.1579 * \text{age} + 2.5797$, $R^2 = 0.39$ ($t_{15} = 3.02$, $P = 0.009$) (Fig. 1). We sampled 4 black-legged kittiwake chicks aged of 10, 14, 40, and 44 days. Their pentosidine concentrations in collagen ranged from $1.84\ \text{pmol}/\text{mg}$ to $10.45\ \text{pmol}/\text{mg}$ (Fig. 1). Like in the adult kittiwakes, the pentosidine levels increased linearly with age: pentosidine = $0.1642 * \text{age} + 0.2621$, $R^2 = 0.54$ ($t_3 = 1.52$, $P = 0.27$). When we pooled adults and chicks, pentosidine levels were independent of age (age: $t_{31} = -0.05$, $P = 0.96$; age²: $t_{31} = 0.49$, $P = 0.63$; sex: $t_{31} = 0.44$, $P = 0.66$; $R^2 = 0.06$). However, if we removed one chick, the relationship improved considerably (age: $t_{30} = 3.07$, $P = 0.004$; age²: $t_{30} = -0.82$, $P = 0.42$; sex: $t_{30} = 0.72$, $P = 0.74$; $R^2 = 0.23$). Collagen content from the standardized 6 mm skin biopsy increased with age in chicks (albeit not significantly) but not in adults (chicks: age: $t_3 = 3.01$, $P = 0.09$; age²: $t_2 = 0.14$, $P = 0.91$; adults: age: $t_{27} = 0.68$, $P = 0.50$; age²: $t_{27} = -0.48$, $P = 0.63$; sex: $t_{27} = 0.54$, $P = 0.60$; Fig. 2).




Fig. 1. Pentosidine/collagen (pmol/mg) relative to age in (a) black-legged kittiwake chicks, (b) black-legged kittiwake adults, (c) auk chicks (puffins and razorbills) and (d) auk adults (puffins and murre; includes murre chicks of unknown exact age).

3.2. Auks

We sampled 14 adult puffins aged 5–26 years and 15 chicks aged 15 to > 35 days old. All age groups were represented by one adult, but no chicks aged 19, 20, 22, 23, 27, 31, 32, and 34 days were captured. Pentosidine concentrations did not vary with age in adults ($R^2 = 0.02$, age: $t_{12} = 0.47$, $P = 0.65$; age²: $t_{12} = 0.44$, $P = 0.67$; Fig. 1). Concentrations decreased linearly with age in the chicks: pentosidine = $-0.1545 * \text{age} + 7.3278$ ($R^2 = 0.39$, age: $t_{14} = -2.91$, $P = 0.01$; age²: $t_{12} = -2.17$, $P = 0.05$). There was no relationship for pooled chick and adult data ($R^2 = 0.17$, age: $t_{42} = 1.43$, $P = 0.16$; age²: $t_{42} = -0.64$, $P = 0.52$). Collagen content of 6 mm biopsies increased with age in chicks but not adults (chicks: age: $t_{13} = 3.81$, $P = 0.002$; age²: $t_{12} = 1.08$, $P = 0.30$; adults: age: $t_{11} = 0.8$, $P = 0.42$; age²: $t_{11} = -0.80$, $P = 0.44$; Fig. 2). We sampled 8 razorbill chicks aged 3–15 days and pentosidine concentrations did not vary with age ($R^2 = 0.08$; age: $t_7 = 0.74$, $P = 0.49$; age²: $t_7 = 0.55$, $P = 0.61$). Collagen increased with age, albeit not significantly (age: $t_6 = 1.61$, $P = 0.16$; age²: $t_5 = 0.65$, $P = 0.54$; Fig. 2). We sampled 5 thick-billed murre chicks of unknown ages (< 8 d old) and 37 adults between 6 and 32 years of age. For pooled data (chicks and adults), pentosidine concentrations did not vary with age ($R^2 = 0.04$, age: $t_{43} = -1.30$, $P = 0.20$; age²: $t_{43} = -1.13$, $P = 0.27$; sex: $t_{43} = 0.70$, $P = 0.49$) (Fig. 1). No trend was apparent for chicks and adults separately (Fig. 1). Collagen was higher in adults than chicks (age: $t_{44} = 2.87$, $P = 0.006$; age²: $t_{43} = -1.44$, $P = 0.16$; Fig. 2).

4. Discussion

Unlike some previous studies on primarily short-lived species (Iqbal et al. 1999a; Chaney et al., 2003; Fallon et al., 2006; Cooley et al., 2010; Dorr et al., 2017), pentosidine concentrations in collagen did not increase substantially with age in the four long-lived seabird species examined in the current study. These results are consistent with the idea that long-lived birds do not show many of the classic traits of physiological aging present in mammals and short-lived birds (Nisbet, 2001; Ricklefs, 2008). Only kittiwakes (the shortest-lived species) showed a significant increase in pentosidine with age in adults, but there was no significant relationship for pooled chick and adults values. Common gulls and bridled terns, which are relatively long-lived, also showed no significant increase with age in pentosidine measured in biopsies from the wing patagium (Rattiste et al., 2015; Labbé et al., 2019), although California gull foot webs showed an increase (Chaney et al., 2003). The California gull study demonstrated lower pentosidine levels in the foot webs of young (~4 years) than old (~20 years) gulls, but showed no change with age within either age class. Without information from intervening age classes, such trends may represent environmental variation associated with each cohort. Skin collagen half-life is about 15 years (Iverson et al., 2017), and consequently the pentosidine technique may be less reliable in long-lived species, as implied by recent work on turtles (Iverson et al., 2017). Moreover, the foot web is more exposed to ultraviolet radiation than the breast, which could lead to degradation of pentosidine, and so levels are likely lower than other skin sites (as is the case for the patagium, Cooley et al., 2010) and trends




Fig. 2. Collagen content (g collagen/L of buffer) relative to age in (a) black-legged kittiwake chicks, (b) black-legged kittiwake adults, (c) auk chicks (puffins and razorbills) and (d) auk adults (puffins and murre chicks of unknown exact age). Values are reported as hydroxyproline content from a 6 mm punch biopsy from the foot webbing of each individual (without accounting for tissue mass), which was converted to collagen by dividing by 0.14.

with age may be weaker. Collagen per sample increased with age, likely because the foot webs became thicker with age, and so more tissue was collected per sample of constant diameter.

We argue that pentosidine is unlikely to be a useful biomarker for chronological age in the foot web of birds. Iverson et al. (2017) found that “concentrations were too variable to permit precise estimates of age for turtles of unknown age” and the same appears to be true for volant vertebrates. The earlier successful papers (on breast or patagial skin tissue) primarily demonstrated an increase in pentosidine with age during the first few years of life. Indeed, our own data showed a relationship between pentosidine and age during chick development, which would be unexpected for a metric that is purported to provide age over an entire animal’s lifespan. Chaney et al. (2003) showed higher pentosidine in California gulls aged 200–300 months than those aged about 50 months. Given that reproductive maturity is typically 48 months, this comparison was between gulls breeding for the first time and those that had bred for many years. Similarly, the increase in monk parakeets was primarily between those aged 6 months or less and those aged 20 months or more (Cooley et al., 2010) and the increase in double-crested cormorants was primarily between those younger than 50 months and those older than 50 months (Fallon et al., 2006; Cooley, 2008). Given that moult patterns can distinguish birds in their first year or two of reproduction, there seems little need for pentosidine to make such distinctions. Moreover, within a single family (gulls, Laridae), the patterns included no variation with age (Rattiste et al., 2015), an increase between young breeders and old breeders (Chaney et al., 2003)

and no change between chicks and adults, but a weak increase with age in breeders (our study). Taken together, our results combined with the literature suggests that this technique does not appear to be reliable for precise aging of wild seabirds. This is likely especially true for foot web biopsies because foot web biopsies have high levels of keratin and low levels of collagen, so pentosidine levels are relatively low.

Kittiwakes were the only species that showed a significant trend in adults. Pentosidine also increased with age in chicks, but the relationship was driven by a single outlier that had the highest level recorded in our study. Indeed, without that outlier, there was a significant increase from chicks to adults that followed accurately the trend derived among adult birds. When averaged across age groups (each year), the r^2 value increased to 0.39, implying that the technique may have potential utility for obtaining an approximate age in this species.

The puffin data was from a poor year when most nests failed; consequently, our sample sizes were small and we had low statistical power. As was the case for kittiwakes, the maximum value for puffin chicks was higher than the maximum value for adults, which challenges the idea that pentosidine increases regularly with age. Nonetheless, our values may be biased because those individuals that were starving or that had starved (and were therefore not sampled) might have had low glucose and therefore less AGEs (Iqbal et al. 1999a). Similarly, our sample size for razorbills was also likely too small to make general conclusions, as we were unable to catch any adults (all chicks died). For mures, we did not know the age of the chicks, but they were no > 10 days old. Interestingly, the lowest level of pentosidine was in birds

aged 6–7 years (due to high collagen, Fig. 2), and values then increased. Again, this is consistent with the idea that pentosidine is lower in birds just starting to breed compared to those breeding later.

Previous studies that found relationships between age and pentosidine used only patagium or breast tissue from necropsied or euthanized birds (Iqbal et al. 1999a; Chaney et al., 2003; Fallon et al., 2006; Cooley et al., 2010). These studies included both wild and domestic birds, and so included both presumably stressed and unstressed individuals. Thus, although food restriction (reductions in glucose and other nutrition) is known to alter pentosidine expression in birds (Iqbal et al. 1997, 1999a), the differences among studies is unlikely to be explained by nutritional stress. The only other studies that obtained samples from live birds found no relationship (Le Souëf, 2012; Rattiste et al., 2015; Campion, 2015; Labbé et al., 2019). Despite high statistical power to detect an effect (e.g., 87% for kittiwakes), we did not detect such a relationship. One possibility is that glucose may react differently post-death, creating altered levels of pentosidine, thus explaining why trends were found for birds that were already dead, but not for live birds (Pozhitkov et al., 2017).

Aging in wild, long-lived birds is likely multifactorial, and there is no single compound that appears to explain a significant proportion of senescence (Ricklefs, 2008; Monaghan et al., 2008; Munshi-South and Wilkinson, 2010). For example, whereas telomere length changes consistently with age in short-lived birds, no such pattern occurs in long-lived birds (Haussmann et al., 2003; Young et al., 2013; Rattiste et al., 2015). For long-lived birds, 'biological' rather than chronological age may be important, and physiological aging may be a complicated combination of multiple factors that change with time (i.e., breeding state, glucose in diet, stress; Ricklefs, 2008; Monaghan and Haussmann, 2006). Additional work, especially longitudinally (sampling the same individual over their lifespans), could address these issues, as well as also address the issue of selective mortality. That is, birds with lower pentosidine might die young, and so the increase in pentosidine with age may be simply an artefact of a cross-sectional sampling design.

CRedit authorship contribution statement

Angelika A. Aleksieva: Investigation. **Jason R. Treberg:** Supervision. **Antony W. Diamond:** Project administration. **Scott A. Hatch:** Project administration. **Kyle H. Elliott:** Supervision.

Acknowledgements

We would like to thank NSERC Discovery Grants to KHE and JT, which helped pay for assays. We would also like to thank a Mobility Award from McGill University, which paid for AA's travel from Montreal to Winnipeg for assays. J. Cunningham, S. Kahane-Rapport and E. Love-Kernick helped in the field at Middleton. The Middleton Island and Coats Island field seasons were paid for by the Canada Research Chairs in Arctic Ecology, the Fonds de Recherche du Québec Nature & Technologie and the McGill University start-up fund. The lab assays were partially paid for by the Canada Research Chair in Environmental Dynamics and Metabolism. This chapter was part of AA's Master's, and we thank J. Head for her help as committee member. Work on Machias Seal Island and Coats Island was supported by Environment and Climate Change Canada. We thank all of the many banders who helped create these known-age populations, especially T. Gaston without whose foresight and decades of dedication the known-age Coats population would not be possible.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.exger.2020.110833>.

References

- Apanius, V., Nisbet, I.C., 2006. Serum immunoglobulin G levels are positively related to reproductive performance in a long-lived seabird, the common tern (*Sterna hirundo*). *Oecologia* 147, 12–23.
- Bagenal, T.B., 1974. *The Ageing of Fish*. Unwin Brothers, Old Woking.
- Broughton, J.M., Rampton, D., Holanda, K., 2002. A test of an osteologically based age determination technique in the double-crested cormorant *Phalacrocorax auritus*. *Ibis* 144, 143–146.
- Brown, K., Jimenez G., A., Whelan, S., Lalla, K., Hatch, A., S., Elliott H., K., 2019. Brown, K., Jimenez, A.G., Whelan, S., Lalla, K., Hatch, S.A. and Elliott, K.H., Muscle fiber structure in an aging long-lived seabird, the black-legged kittiwake (*Rissa tridactyla*). *Journal of Morphology* 280 (7), 1061–1070.
- Campion, M.R., 2015. The simultaneous determination of hydroxyproline and pentosidine in avian skin tissue. *Chem. New Zealand* 79, 172–176.
- Chaney, R.C., Blemings, K.P., Bonner, J., Klandorf, H., 2003. Pentosidine as a measure of chronological age in wild birds. *Auk* 120, 394–399.
- Chivers, L.S., Hatch, S.A., Elliott, K.H., 2015. Accelerometry reveals an impact of short-term tagging on seabird activity budgets. *Condor* 118, 159–168.
- Cichon, M., Sendek, J., Gustafsson, L., 2003. Age-related decline in humoral immune function in collared flycatchers. *J. Evol. Biol.* 16, 1205–1210.
- Cooley, C.K., 2008. Development and Evaluation of a Minimally Invasive Sampling Technique to Estimate the Age of Living Birds. MSc thesis. West Virginia University.
- Cooley, C.K., Fallon, J.A., Avery, M.L., Anderson, J.T., Falkenstein, E.A., Klandorf, H., 2010. Refinement of biomarker pentosidine methodology for use on aging birds. *Human-Wildlife Interactions* 4, 304–314.
- Coulson, J.C., Fairweather, J.A., 2003. Reduced reproductive performance prior to death in the black-legged kittiwake: senescence or terminal illness? *J. Avian Biol.* 32, 146–152.
- Daffu, G., del Pozo, C., O'Shea, K., Ananthakrishnan, R., Ramasamy, R., Schmidt, A., 2013. Radical roles for RAGE in the pathogenesis of oxidative stress in cardiovascular diseases and beyond. *Int. J. Mol. Sci.* 14, 19891–19910.
- Dammann, P., Sell, D.R., Begall, S., Strauch, C., Monnier, V.M., 2011. Advanced glycation end-products as markers of aging and longevity in the long-lived Ansell's mole-rat (*Fukomys ansellii*). *J. Gerontol. A, Biomed. Sci. Med. Sci.* 67, 573–583.
- Diamond, A.W., Devlin, C.M., 2003. Seabirds as indicators of changes in marine ecosystems, ecological monitoring on Machias Seal Island. *Environ. Monitoring Assess.* 88, 153–181.
- Dorr, B.S., Stahl, R.S., Hanson-Dorr, K.C., Furcolow, C.A., 2017. Using pentosidine and hydroxyproline to predict age and sex in an avian species. *Ecology and Evolution* 7, 8999–9005.
- Dyer, D.G., Blackledge, J.A., Thorpe, S.R., Baynes, J.W., 1991. Formation of pentosidine during nonenzymatic browning of proteins by glucose. Identification of glucose and other carbohydrates as possible precursors of pentosidine in vivo. *J. Biol. Chem.* 266, 11654–11660.
- Elliott, K.H., O'Reilly, K.M., Hatch, S.A., Gaston, A.J., Hare, J.F., Anderson, W.G., 2014. The prudent parent meets old age, a high stress response in very old seabirds supports the prudent restraint hypothesis. *Horm. Behav.* 66, 828–837.
- Elliott, K.H., Hare, J.F., Le Vaillant, M., Gaston, A.J., Ropert-Coudert, Y., Anderson, W.G., 2015. Ageing gracefully, physiology but not behaviour declines with age in a diving seabird. *Funct. Ecol.* 29, 219–228.
- Fallon, J.A., Cochrane, R.L., Dorr, B., Klandorf, H., 2006. Interspecies comparison of pentosidine accumulation and its correlation with age in birds. *Auk* 123, 870–876.
- Gaston, A.J., de Forest, L.N., Donaldson, G., Noble, D.G., 1994. Population parameters of thick-billed murrelets at Coats Island, Northwest Territories, Canada. *Condor* 96, 935–948.
- Gill, V.A., Hatch, S.A., 2002. Components of productivity in black-legged kittiwakes *Rissa tridactyla*, response to supplemental feeding. *J. Avian Biol.* 33, 113–126.
- Grandhee, S.K., Monnier, V.M., 1991. Mechanism of formation of the Maillard protein cross-link pentosidine. *J. Biol. Chem.* 266, 11649–11653.
- Grue, H., Jensen, B., 1979. Review of the formation of incremental lines in tooth cementum of terrestrial mammals. *Danish Rev. Game Biol.* 11, 1–48.
- Harman, D., 1992. Free radical theory of aging. *Mutation Res. DNAGing* 275, 257–266.
- Harman, D., 2009. Origin and evolution of the free radical theory of aging, a brief personal history, 1954–2009. *Biogerontol* 10, 773–781.
- Haussmann, M.F., Winkler, D.W., O'Reilly, K.M., Huntington, C.E., Nisbet, I.C., Vleck, C.M., 2003. Telomeres shorten more slowly in long-lived birds and mammals than in short-lived ones. *Proc. Roy. Soc. Lond. B* 270, 1387–1392.
- Herman, D.P., Ylitalo, G.M., Robbins, J., Straley, J.M., Gabriele, C.M., Clapham, P.J., Boyer, R.H., Tilbury, K.L., Pearce, R.W., Krahn, M.M., 2009. Age determination of humpback whales *Megaptera novaeangliae* through blubber fatty acid compositions of biopsy samples. *Mar. Ecol. Prog. Ser.* 392, 277–293.
- Holmes, D., Martin, K., 2009. A bird's-eye view of aging, what's in it for ornithologists? *Auk* 126, 1–23.
- Holmes, D.J., Ottinger, M.A., 2003. Birds as long-lived animal models for the study of aging. *Exp. Gerontol.* 38, 1365–1375.
- Iqbal, M., Probert, L.L., Klandorf, H., 1997. Effect of dietary aminoguanidine on tissue pentosidine and reproductive performance in broiler breeder hens. *Poultry Sci* 76, 1574–1579.
- Iqbal, M., Kenney, P.B., Klandorf, H., 1999a. Age-related changes in meat tenderness and tissue pentosidine, effect of diet restriction and aminoguanidine in broiler breeder hens. *Poultry Sci* 78, 1328–1333.
- Iqbal, M., Probert, L.L., Alhumadi, N.H., Klandorf, H., 1999b. Protein glycosylation and advanced glycosylated endproducts (AGEs) accumulation, an avian solution? *J. Gerontol. Biol. Sci.* 54, B171–B176.

- Iverson, J.B., Stahl, R.S., Furcolow, C., Kraus, F., 2017. An evaluation of the use of pentosidine as a biomarker for ageing turtles. *Conserv. Physiol.* 5 (76).
- Jarman, S.N., Polanowski, A.M., Faux, C.E., Robbins, J., de Paoli-Iseppi, R., Bravington, M., Deagle, B.E., 2015. Molecular biomarkers for chronological age in animal ecology. *Mol. Ecol.* 24, 4826–4847.
- Jimenez G., A., O'Connor S., E., Elliott H., K., 2019. Muscle myonuclear domain, but not oxidative stress, decreases with age in a long-lived seabird with high activity costs. *Journal of Experimental Biology* 222 (18), jeb211185.
- Jones, O.R., Scheuerlein, A., Salguero-Gómez, R., Camarda, C.G., Schaible, R., Casper, B.B., Dahlgren, J.P., Ehrlén, J., García, M.B., Menges, E.S., Quintana-Ascencio, P.F., 2014. Diversity of ageing across the tree of life. *Nature* 505, 169.
- Kelly, K.G., Diamond, A.W., Holberton, R.L., Bowser, A.K., 2015. Researcher handling of incubating Atlantic puffins *Fratercula arctica* has no effect on reproductive success. *Mar. Ornithol.* 43, 77–82.
- Klomp, N.I., Furness, R.W., 1992. A technique which may allow accurate determination of the age of adult birds. *Ibis* 134, 245–249.
- Labbé, A.M.T., Le Souef, A.T., Dunlop, J.N., Calver, M., Shephard, J.M., van Keulen, M., 2019. Pentosidine levels cannot be used to age a long-lived seabird. *Emu* 119, 186–190.
- Le Souëf, A., 2012. Development and Improvement of Clinical Tools for Rehabilitating Endangered Black Cockatoos (*Calyptorhynchus spp.*) Back to the Wild. Dissertation. Murdoch University, Australia.
- Lecomte, V.J., Sorci, G., Cornet, S., Jaeger, A., Faivre, B., Arnoux, E., et al., 2010. Patterns of aging in the long-lived wandering albatross. *Proc. Natl. Acad. Sci. U. S. A.* 107, 6370–6375.
- Lincoln, F.C., 1921. The history and purposes of bird banding. *Auk* 38, 217–228.
- Lushchak, V.I., 2014. Free radicals, reactive oxygen species, oxidative stress and its classification. *Chem. Biol. Interactions* 224, 164–175.
- Maekawa, T., Rathinasamy, T.K., Altman, K.I., Forbes, W.F., 1970. Changes in collagen with age—I, the extraction of acid soluble collagens from the skin of mice. *Exp. Gerontol.* 5, 177–186.
- Martin, G.M., Austad, S.N., Johnson, T.E., 1996. Genetic analysis of ageing, role of oxidative damage and environmental stresses. *Nat. Gen.* 13, 25–34.
- McNeil, R., Burton, J., 1972. Cranial pneumatization patterns and bursa of fabricius in North American shorebirds. *Wilson Bull.* 84, 329–339.
- Moe, B., Rønning, B., Verhulst, S., Bech, C., 2008. Metabolic ageing in individual zebra finches. *Biology Letters* 5 (1), 86–89.
- Monaghan, P., Haussmann, M.F., 2006. Do telomere dynamics link lifestyle and lifespan? *Trends Ecol. Evol.* 21, 47–53.
- Monaghan, P., Charmantier, A., Nussey, D.H., Ricklefs, R.E., 2008. The evolutionary ecology of senescence. *Funct. Ecol.* 22, 371–378.
- Monaghan, P., Metcalfe, N.B., Torres, R., 2009. Oxidative stress as a mediator of life history trade-offs, mechanisms, measurements and interpretation. *Ecol. Lett.* 12, 75–92.
- Monnier, V.M., Cerami, A., 1981. Nonenzymatic browning in vivo, possible process for aging of long-lived proteins. *Science* 211, 491–493.
- Morris, P.A., 1972. A review of mammalian age determination methods. *Mamm. Rev.* 2, 69–104.
- Munshi-South, J., Wilkinson, G.S., 2010. Bats and birds, exceptional longevity despite high metabolic rates. *Ageing Res. Rev.* 9, 12–19.
- Nisbet, I.C.T., 2001. Detecting and measuring senescence in wild birds: experience with long-lived seabirds. *Exp. Gerontol.* 36, 833–843.
- Nisbet, I.C.T., Apanius, V., Friar, M.S., 2002. Breeding performance of very old common terns. *J. Field Ornithol.* 73, 117–124.
- Nitti, M., d'Abramo, C., Traverso, N., Verzola, D., Garibotto, G., Poggi, A., Odetti, P., Cottalasso, D., Marinari, U.M., Pronzato, M.A., Domenicotti, C., 2005. Central role of PKC δ in glycooxidation-dependent apoptosis of human neurons. *Free Radic. Biol. Med.* 38, 846–856.
- Nowotny, K., Jung, T., Höhn, A., Weber, D., Grune, T., 2015. Advanced glycation end products and oxidative stress in type 2 diabetes mellitus. *Biomolecules* 5, 194–222.
- Nussey, D.H., Coulson, T., Festa-Bianchet, M., Gaillard, J.M., 2008. Measuring senescence in wild animal populations, towards a longitudinal approach. *Funct. Ecol.* 22, 393–406.
- Palacios, M.G., Cunnick, J.E., Winkler, D.W., Vleck, C.M., 2007. Immunosenescence in some but not all immune components in a free-living vertebrate, the tree swallow. *Proc. Roy. Soc. Lond. B* 274, 951–957.
- Peabody, F.E., 1961. Annual growth zones in living and fossil vertebrates. *J. Morphol.* 108, 11–62.
- Pozhitkov, A.E., Neme, R., Domazet-Loso, T., Leroux, B.G., Soni, S., Tautz, D., Noble, P.A., 2017. Tracing the dynamics of gene transcripts after organismal death. *Open Biol.* 7, 1–25.
- Rattiste, K., Klandorf, H., Urvik, J., Sepp, T., Asghar, M., Hasselquist, D., Cooley, C., Hörak, P., 2015. Skin pentosidine and telomere length do not covary with age in a long-lived seabird. *Biogerontol* 16, 435–441.
- Reid, R.E., 1981. Lamellar-zonal bone with zones of annuli in the pelvis of a sauropod dinosaur. *Nature* 292, 49–51.
- Ricklefs, R.E., 2008. The evolution of senescence from a comparative perspective. *Funct. Ecol.* 22, 379–392.
- Ricklefs, R.E., 2010. Insights from comparative analyses of aging in birds and mammals. *Aging Cell* 9, 273–284.
- Rodway, M.S., Montevecchi, W.A., Chardine, J.W., 1996. Effects of investigator disturbance on breeding success of Atlantic Puffins *Fratercula arctica*. *Biol. Conserv.* 76, 311–319.
- Saino, N., Ferrari, R.P., Romano, M., Rubolini, D., Møller, A.P., 2003. Humoral immune response in relation to senescence, sex and sexual ornamentation in the barn swallow (*Hirundo rustica*). *J. Evol. Biol.* 16, 1127–1134.
- Sell, D.R., Monnier, V.M., 1989. Structure elucidation of a senescence cross-link from human extracellular matrix. *J. Biol. Chem.* 264, 21597–21602.
- Sell, D.R., Prime, M., Schafer, I.A., Kovach, M., Weiss, M.A., Monnier, V.M., 1998. Cell-associated pentosidine as a marker of aging in human diploid cells in vitro and in vivo. *Mech. Ageing Develop* 221–240.
- Sell, D.R., Lane, M.A., Obrenovich, M.E., Mattison, J.A., Handy, A., Ingram, D.K., Cutler, R.G., Roth, G.S., Monnier, V.M., 2003. The effect of caloric restriction on glycation and glycooxidation in skin collagen of nonhuman primates. *J. Gerontol. A, Biol. Sci. Med. Sci.* 58, B508–B516.
- Siegel-Causey, D., 1989. Cranial pneumatization in the Phalacrocoracidae. *Condor* 90, 885–905.
- Sugimori, F., Oka, N., Ishibashi, Y., 1985. The degree of skull ossification as a means of aging short-tailed shearwaters. *J. Yamashina Inst. Ornithol.* 17, 159–165.
- Turbill, C., Ruf, T., 2010. Senescence is more important in the natural lives of long- than short-lived mammals. *PLoS One* 5, e12019.
- Valko, M., Leibfritz, D., Moncol, J., Croning, M.T.D., Mazur, M., Telser, J., 2007. Free radicals and antioxidants in normal physiological functions and human disease. *The International J. Biochem. Cell Biol.* 39, 44–84.
- Wickens, A.P., 2001. Ageing and the free radical theory. *Resp. Physiol.* 128, 379–391.
- Young, R.C., Kitaysky, A.S., Haussmann, M.F., Descamps, S., Orben, R.A., Elliott, K.H., Gaston, A.J., 2013. Age, sex, and telomere dynamics in a long-lived seabird with male-biased parental care. *PLoS One* 8, e74931.